

Descent Polynomials, Peak Polynomials and an Involution on Permutations

Ezgi Kantarcı Oğuz

Department of Mathematics, Sabancı University, Turkey

Abstract. The size of the set of all permutations of n with a given descent set is a polynomial in n , called the descent polynomial. Similarly, the size of the set of all permutations of n with a given peak set, adjusted by a power of 2 gives a polynomial in n , called the peak polynomial. We give a unitary expansion of descent polynomials in terms of peak polynomials. Then we use this expansion, along with an involution that flips the initial segment of a permutation, to give a combinatorial interpretation of the coefficients of the peak polynomial in a binomial basis, thus giving a new proof of the peak polynomial positivity conjecture.

Keywords: permutation, descent, peak, spike

1 Introduction

Denote by \mathfrak{S}_n the symmetric group of permutations $\sigma = \sigma_1\sigma_2 \cdots \sigma_n$ of $[n] = \{1, 2, \dots, n\}$ written in one-line notation. We will draw the graph of σ by plotting points (i, σ_i) and connecting consecutive points.

We define the descent set of σ as follows:

$$\text{Des}(\sigma) = \{i \mid \sigma_i > \sigma_{i+1}\} \subset [n-1].$$

Note that the descents mark the beginnings of the intervals where the graph is decreasing, as seen in [Figure 1](#) below.




Figure 1: The graph of $\sigma = 24315678$ with descents marked in blue.

For a given set S and $n > \max(S)$, we let $D(S, n)$ be the set of all permutations in \mathfrak{S}_n with descent set S , and put $d(S, n) = |D(S, n)|$. In 1915, it was shown by MacMahon [4] that this is a polynomial in n . More recently, Diaz-Lopez et al. [3] proved this polynomial

expands into the binomial basis around $n - m$, where $m = \max(S)$, and gave a combinatorial interpretation for the coefficients. Using the notation $\sigma|_i = \{\sigma_1, \sigma_2, \dots, \sigma_i\}$, we give a version of their result slightly altered to include the case $m > \max(S)$ as follows:

Theorem 1.1 ([3]). *For any finite set of positive integers S with $\max(S) \leq m$ we have:*

$$d(S, n) = a_0(S) \binom{n-m}{0} + a_1(S) \binom{n-m}{1} + \dots + a_m(S) \binom{n-m}{m}, \quad (1.1)$$

where the constant $a_k(S)$ is the number of $\sigma \in D(S, 2m)$ such that:

$$\sigma|_m \cap [m+1, 2m] = [m+1, m+k].$$

The underlying idea is quite elegant and will be useful when we are proving a similar result for peak polynomials. Simply put, if $\sigma \in D(S, n)$ has $\{\sigma_1, \sigma_2, \dots, \sigma_m\} \cap [m+1, n] = A$ for some k element set A , and B is any other k element subset of $[m+1, n]$, then exchanging elements of A and B while preserving the orders gives another permutation with descent set S . Therefore, for each k , it is enough to count for the simplest k -element subset of $[m+1, n]$ and multiply with $\binom{n-m}{k}$.

For example, there are 3 elements in $D(\{2, 3\}, n)$ satisfying $\sigma|_m \cap [5, 8] = \emptyset$: 14325678, 24315678, 34215678. So we have $a_0(\{2, 3\}) = 3$ for $m = 4$. Calculating the other coefficients similarly, we obtain:

$$d(\{2, 3\}, n) = 3 \binom{n-4}{0} + 8 \binom{n-4}{1} + 7 \binom{n-4}{2} + 2 \binom{n-4}{3} + 0 \binom{n-4}{4}. \quad (1.2)$$

Another well studied permutation statistic is given by peak point. Here we define the peak points and their counterpart valley points of the partition to be the points higher and lower than their neighbors respectively:

$$\text{Peak}(\sigma) = \{\sigma \mid \sigma_i > \sigma_{i+1}, \sigma_{i-1}\} \subset [n-1]/\{1\},$$

$$\text{Valley}(\sigma) = \{\sigma \mid \sigma_i < \sigma_{i+1}, \sigma_{i-1}\} \subset [n-1]/\{1\}.$$

The example $\sigma = 34215678$ from **Figure 1** has $\text{Peak}(\sigma) = \{2\}$ and $\text{Valley}(\sigma) = \{4\}$. We also set $\text{Spike}(\sigma) = \text{Peak}(\sigma) \cup \text{Valley}(\sigma)$ to be the set of all extremal points that are not corner points.

For a given set I and $n > \max(I)$, we let $P(I, n)$ be the set of permutations in \mathfrak{S}_n with peak set I , and set $p(I, n) = 2^{-n+|I|+1} |P(I, n)|$. Note that peaks are more restrictive in the sense that $p(I, n) = 0$ if I contains 1 or any consecutive entries. For the rest of this work, we will focus our attention to *admissible* peak sets I : $I \subset [n-1]/\{1\}$ such that $i \in I \Rightarrow i+1 \notin I$.

In [1] Billey, Burzdy and Sagan proved that $p(I, n)$ is a polynomial in n , and conjectured that the coefficients of this polynomial in a binomial basis centered at $\max(I)$ are

non-negative. Their conjecture was proved in 2017 by Diaz-Lopez et al. [2] using the recursion of peaks, without describing the actual coefficients.

In this work, we tie the theory of peak and descent polynomials together by giving a binary expansion of $d(S, n)$ in terms of peak polynomials. We use this expansion to give a description of the peak polynomial coefficients analogous to the one in [Theorem 1.1](#). In [Section 2](#), we extend our notions of descents and peaks to \mathfrak{B}_n , the set of marked permutations of n with $2^n n!$ elements. The added exponent of 2 cancels out with the $2^{-n+|I|+1}$ from the peak polynomial definition, giving us a way to expand descent polynomials in terms of peak polynomials. In [Section 3](#), we define involutions on permutations that flip the descents on an initial section and we use them to partition permutations with a given descent set to calculate the coefficients for the peak polynomial.

2 Descents and Peaks of Marked Permutations

We start our section by tweaking our notation a little bit to express our formulas easier. Note that the peaks and valleys of a permutation only depend on its descent set. In fact for any $S \in [n-1]$, we can talk about the peaks and valleys of S :

$$\text{Peak}(S) = \{1 < i \leq n-1 \mid i \in S, i-1 \notin S\},$$

$$\text{Valley}(S) = \{1 < i \leq n-1 \mid i \notin S, i-1 \in S\}.$$

Note that with this notation, $\text{Peak}(\sigma) = \text{Peak}(\text{Des}(\sigma))$ and $\text{Valley}(\sigma) = \text{Valley}(\text{Des}(\sigma))$ as expected. We also set $\text{Spike}(S) := \text{Peak}(S) \cup \text{Valley}(S)$.

Denote by \mathfrak{B}_n the set of signed permutations:

$$\mathfrak{B}_n := \{\rho = \rho_1 \rho_2 \dots \rho_n \mid \forall i \leq n \exists k : \rho_k = i \text{ or } \rho_k = -i\}.$$

Note that the definitions of descent, peak, spike and valley naturally extend to signed permutations by saying i is a descent of ρ if $\rho_i > \rho_{i+1}$.

Lemma 2.1. *Let $\sigma \in S_n$ have $\text{Peak}(\sigma) = I$. Denote by $\text{Mark}(\sigma)$ the 2^n element subset of \mathfrak{B}_n that give σ when marks are erased. Then, for all $\rho \in \text{Mark}(\sigma)$, $\text{Spike}(\rho) \supset I$. Conversely, for any $S \subset [n-1]$ satisfying $\text{Spike}(S) \supset I$, there are exactly $2^{|I|+1}$ elements in $\text{Mark}(\sigma)$ with descent set equal to S .*

Theorem 2.2. *We have $d(S, n) = \sum_{I \subset \text{Spike}(S)} p(I, n)$.*

For example:

- $d(\emptyset, n) = p(\emptyset, n) = 1$.
- $d(\{1\}, n) = p(\{2\}, n) + p(\emptyset, n)$.




Figure 2: For $I = \{2, 4\}$, $S_I = \{2, 3\}$ as seen above.

- For $1 < k < n$, $d(\{k\}, n) = p(\{k\}, n) + p(\{k+1\}, n) + p(\emptyset, n)$.
- $d(\{k, k+1, \dots, k+j\}, n) = p(\{k, k+j+1\}, n) + p(\{k\}, n) + p(\{k+j+1\}, n) + p(\emptyset, n)$.

For any set $I \in [n]/\{1\}$, we will let S_I denote the unique subset of $[n]/\{n\}$ satisfying $\text{Spike}(S_I) = I$, constructed by alternating the elements of I to be peaks and valleys such that the rightmost one is not a peak. For example, for $I = \{2, 4\}$ we have $S_I = \{2, 3\}$: the descent set with a peak at 2 and a valley at 4 (see [Figure 2](#)).

Corollary 2.3. For any admissible set I , $p(I, n) = \sum_{J \subset I} (-1)^{|I|-|J|} d(S_J, n)$.

If we consider our running example $I = \{2, 4\}$, we get the following formulas from [Theorem 2.2](#) and [Corollary 2.3](#) respectively:

$$d(\{2, 3\}, n) = p(\{2, 4\}, n) + p(\{2\}, n) + p(\{4\}, n) + p(\emptyset, n), \quad (2.1)$$

$$p(\{2, 4\}, n) = d(\{2, 3\}, n) - d(\{1\}, n) - d(\{1, 2, 3\}, n) + d(\emptyset, n). \quad (2.2)$$

3 A combinatorial expression for peak coefficients

We start with defining an operation on permutations that ‘flips’ the orders of some initial coordinates.

Definition 3.1. Let $\sigma \in S_n$. Let $i \leq n$, and $\sigma|_i = \{a_1 < a_2 < \dots < a_i\}$. We define the involution fl_i as follows:

$$\text{fl}_i(\sigma)_j = \begin{cases} a_{i-k+1} & j \leq i, \sigma_j = a_k \\ \sigma_j & j > i. \end{cases}$$

See [Figure 3](#) for examples.

Remark 3.2. The involution fl_i satisfies the following:

- $\text{fl}_i(\sigma)|_i = \sigma|_i$.
- For $k < i$, k is a descent of $\text{fl}_i(\sigma)$ iff it is not a descent of σ .




Figure 3: Operations fl_2 and fl_4 on $\sigma = 24315678$.

- For $k > i$, k is a descent of $fl_i(\sigma)$ iff it is a descent of σ .
- fl_i exchanges all the peaks less than i with valleys, and all the valleys less than i with peaks.

In Figure 3, we see an instance of operations fl_2 and fl_4 commuting. Now we will prove that this is the case in general.

Proposition 3.3. For all i, j , fl_i and fl_j commute.

Definition 3.4. For $i \in \text{Spike}(\sigma)$, σ is said to admit an i^+ -flip if $\text{Spike}(fl_i(\sigma)) = \text{Spike}(\sigma) / \{i\}$. Similarly, it is said to admit an i^- -flip if $\text{Spike}(fl_{i-1}(\sigma)) = \text{Spike}(\sigma) / \{i\}$. We say σ admits an i -flip if it admits an i^+ - or i^- -flip. For σ that admits an i -flip, we set

$$\Psi_i(\sigma) = \begin{cases} fl_i(\sigma) & \sigma \text{ admits an } i^+ \text{-flip,} \\ fl_{i-1}(\sigma) & \text{otherwise.} \end{cases}$$

Visually, this means that fl_i or fl_{i-1} straightens out the peak or valley point at i . For example, 24315678 from Figure 3 admits a 4-flip, but not a 2-flip.

Proposition 3.5. For all i, j such that $|i - j| > 1$, σ admits an i^+ -flip if and only if $\Psi_j(\sigma)$ admits an i^+ -flip. Similarly, σ admits an i^- -flip if and only if $\Psi_j(\sigma)$ admits an i^- -flip.

For any admissible set $I = \{i_1, i_2, \dots, i_k\}$, we put $\Psi_I := \Psi_{i_1} \circ \Psi_{i_2} \circ \dots \circ \Psi_{i_k}$. This operation is well-defined by Proposition 3.3, and the corollary above.

Lemma 3.6. For any admissible set I and any $J \subset I$, Ψ_J induces a bijection between elements of $D(S_I, n)$ that admit a j -flip for all $j \in J$ and elements of $D(S_{I/J}, n)$. In particular, for any $m \geq \max(I)$ we have:

$$\sigma|_m \cap [m + 1, 2m] = [m + 1, m + k] \iff \Psi_J(\sigma)|_m \cap [m + 1, 2m] = [m + 1, m + k].$$

Theorem 3.7. For any admissible set of I with $\max(I) \leq m$ we have

$$p(I, n) = b_0(I) \binom{n-m}{0} + b_1(I) \binom{n-m}{1} + \cdots + b_m(I) \binom{n-m}{m}, \quad (3.1)$$

where the constant $b_k(I)$ is the number of $\sigma \in D(S_I, 2m)$ such that:

$$\sigma|_m \cap [m+1, 2m] = [m+1, m+k],$$

and σ does not admit any i -flips.

	2-flip	4-flip		2-flip	4-flip		2-flip	4-flip
14325678	✗	✓	15324678	✗	✓	16523478	✗	✗
24315678	✗	✓	15423678	✗	✗	26513478	✗	✗
34215678	✓	✓	25314678	✗	✗	36512478	✗	✗
			25413678	✗	✗	46512378	✗	✗
			35214678	✓	✗	56213478	✓	✗
57612348	✗	✗	35412678	✗	✗	56312478	✓	✗
675123478	✓	✗	45213678	✓	✗	56412378	✓	✗
			45312678	✓	✗			
	$k = 0, k = 3$			$k = 1$			$k = 2$	

Table 1: The elements $\sigma \in D(\{2, 3\}, 8)$ satisfying $\sigma|_4 \cap [5, 8] = [5, 4+k]$.

We will end this section by calculating the expansion of $p(\{2, 4\}, n)$. Recall that $S_{\{2,4\}} = \{2, 3\}$. For all elements $\sigma \in D(\{2, 3\}, 8)$ satisfying $\sigma|_4 \cap [5, 8] = [5, 4+k]$ for some k we need to check if σ admits a 2-flip or a 4-flip. Checking for 2-flips is very straightforward, we just need to check whether $\sigma_1 > \sigma_3$. 4-flips are slightly more tricky as fl_4 does not simply exchange a pair of coordinates, and we actually need to calculate $\text{fl}_4(\sigma)$ to see if $\text{fl}_4(\sigma)_3$ is smaller than σ_5 . For each k , the related permutations σ can be found in [Table 1](#), along with the information on whether they admit 2 or 4-flips.

Counting the elements that admit neither 2 nor 4-flips from [Table 1](#) gives us the following formula:

$$p(\{2, 4\}, n) = 0 \binom{n-4}{0} + 4 \binom{n-4}{1} + 4 \binom{n-4}{2} + 1 \binom{n-4}{3}.$$

In fact, the inclusion-exclusion principle allows us to read the coefficients for $p(2, n)$ (ones that admit only 4-flips), $p(4, n)$ (ones that admit only 2-flips) and $p(\emptyset, n)$ (ones that admit

only 4-flips) from [Table 1](#):

$$\begin{aligned} p(\{2\}, n) &= 2\binom{n-4}{0} + 1\binom{n-4}{1} + 0\binom{n-4}{2} + 0\binom{n-4}{3}, \\ p(\{4\}, n) &= 0\binom{n-4}{0} + 3\binom{n-4}{1} + 3\binom{n-4}{2} + 1\binom{n-4}{3}, \\ p(\emptyset, n) &= 1\binom{n-4}{0} + 0\binom{n-4}{1} + 0\binom{n-4}{2} + 0\binom{n-4}{3}. \end{aligned}$$

Note that $p(\{2,4\}, n) + p(\{2\}, n) + p(\{4\}, n) + p(\emptyset, n) = d(\{2,3\}, n)$ as required.

Acknowledgements

The author would like to thank Mohamed Omar for an inspiring seminar talk on the subject. The author is also immensely grateful to Alexander Diaz-Lopez and Erik Insko for spotting an error with the initial statement of the main result, and their many helpful suggestions and comments in the following discussion. This work was partially supported by the USC Graduate School Final Year Fellowship.

References

- [1] S. Billey, K. Burdzy, and B. E. Sagan. “Permutations with given peak set”. *J. Integer Seq.* **16.6** (2013), Article 13.6.1, 18 pp. [Link](#).
- [2] A. Diaz-Lopez, P. E. Harris, E. Insko, and M. Omar. “A proof of the peak polynomial positivity conjecture”. *J. Combin. Theory Ser. A* **149** (2017), pp. 21–29. [Link](#).
- [3] A. Diaz-Lopez, P. E. Harris, E. Insko, M. Omar, and B. E. Sagan. “Descent polynomials”. 2017. [arXiv:1710.11033](#).
- [4] P. A. MacMahon. *Combinatory Analysis. Vol. I, II (bound in one volume)*. Dover Phoenix Editions. Reprint of *An Introduction to Combinatory Analysis* (1920) and *Combinatory Analysis. Vol. I, II* (1915, 1916). Dover Publications, Inc., Mineola, NY, 2004.